


How well is the health care system in New York working for women?

New York ranks #10 overall.

Worse-than-average performance



Better-than-average performance

New York ranks

5

of 51

on Health and Reproductive Care Outcomes

Maternal and all-cause women's mortality + Infant mortality + Physical and mental health status

10

of 51

on Coverage, Access, and Affordability

Insurance coverage + Provider accessibility + Health care affordability


27

of 51

on Health Care Quality and Prevention

Low-risk cesarean birth rate + Preventive care use + Pre- and postpartum care + Mental health care screening

New York ranks #2 in the Mid-Atlantic region.



New York performed best on:

- ▲ Women ages 18–64 who have ever had an HIV or AIDS test (3 of 51)
- ▲ Women with depression before or during pregnancy (3 of 33)
- ▲ Self-pay in-hospital births (3 of 51)

New York performed worst on:

- ▼ Women with up-to-date colon cancer screening (34 of 51)
- ▼ Women age 65 and older who have ever received a pneumonia vaccine (43 of 51)
- ▼ Low-risk cesarean births per 100 women giving live birth (48 of 51)

Women's Health and Reproductive Care Indicator Data by Dimension

| Dimension and indicator | Data year | NY rate | U.S. average | Best state rate | State rank ^a |
|---|-------------|---------|--------------|-----------------|-------------------------|
| Health Outcomes | | | | | |
| Maternal deaths while pregnant or within 42 days of termination of pregnancy, per 100,000 live births | 2020 - 2022 | 23.1 | 26.3 | 0.0 | 12 of 43 |
| Infant mortality, deaths per 1,000 live births | 2021 | 4.2 | 5.4 | 2.8 | 9 of 51 |
| Breast and cervical cancer deaths per 100,000 female population | 2022 | 17.3 | 20.9 | 15.3 | 6 of 51 |
| All-cause mortality rate per 100,000 women ages 15–44 | 2022 | 81.9 | 110.3 | 70.5 | 5 of 51 |
| Percent of reported live births where baby was born prior to 37 weeks of pregnancy (gestation), among birth records that reported a gestational age | 2022 | 9% | 10% | 8% | 12 of 51 |
| Rate of women ages 15–44 with syphilis per 100,000 female population | 2022 | 46.0 | 78.0 | 1.0 | 18 of 51 |
| Rate of infants born with congenital syphilis per 100,000 live births | 2022 | 23.7 | 102.5 | 0.0 | 10 of 51 |
| Percent of women ages 18–64 who reported being in fair or poor health | 2022 | 15% | 17% | 11% | 18 of 51 |
| Percent of women with a recent live birth with self-reported depression in the three months before or during pregnancy | 2021 | 16% | 22% | 12% | 3 of 33 |
| Percent of women with a recent live birth with self-reported postpartum depressive symptoms | 2021 | 12% | 13% | 9% | 13 of 33 |
| Percent of women ages 18–64 who reported having 14 or more poor mental health days in the past month | 2022 | 19% | 21% | 15% | 9 of 51 |
| Percent of women with a recent live birth who experienced intimate partner violence before and/or during pregnancy | 2021 | 2.7% | 3.1% | 1.8% | 11 of 33 |
| Coverage, Access, and Affordability | | | | | |
| Percent of women ages 19–64 without health insurance coverage | 2022 | 5% | 10% | 3% | 8 of 51 |
| Percent of women with a recent live birth without health insurance coverage a month before pregnancy | 2021 | 6% | 12% | 2% | 8 of 33 |
| Percent of women with a recent live birth without health insurance coverage during pregnancy | 2021 | 1.2% | 2.6% | .4% | 10 of 33 |
| Percent of women ages 18–44 who reported a time in the past 12 months when they needed to see a doctor but could not because of cost | 2022 | 14% | 17% | 7% | 19 of 51 |
| Percent of women ages 18–44 who did not have one (or more) person they think of as their personal health care provider | 2022 | 21% | 23% | 8% | 31 of 51 |

Women's Health and Reproductive Care Indicator Data by Dimension (cont.)

| Dimension and indicator | Data year | NY rate | U.S. average | Best state rate | State rank |
|--|-----------|---------|--------------|-----------------|------------|
| Coverage, Access, and Affordability (cont.) | | | | | |
| Share of in-hospital births in state with a self-pay insurance payment source | 2022 | .6% | 2.9% | .3% | 3 of 51 |
| Rate of maternity care providers (MDs, DOs, certified nurse midwives practicing in Obstetrics and Gynecology) per 100,000 women ages 15–44 ^b | 2022 | 104.7 | 78.9 | 159.7 | 12 of 51 |
| Abortion clinics per 100,000 women ages 15–44 ^b | 2023 | 2.3 | 1.5 | 12.9 | 19 of 51 |
| Health Care Quality and Prevention | | | | | |
| Rate of singleton, term (37 completed weeks or more of gestation based on the obstetric estimate), vertex (not breech), cesarean deliveries to women having a first birth per 100 women delivering singleton, term, vertex, first births | 2022 | 29.5 | 26.3 | 16.7 | 48 of 51 |
| Percent of women ages 18–44 who reported not visiting a doctor for a routine checkup in the past two years | 2022 | 11% | 13% | 8% | 12 of 51 |
| Percent of live births where prenatal care did not begin during the first to third month of pregnancy, among birth records that specified a time period for when prenatal care began | 2022 | 20% | 23% | 13% | 20 of 51 |
| Percent of women with a recent live birth who did not report receiving a maternal postpartum checkup visit | 2021 | 9% | 9% | 6% | 15 of 33 |
| Percent of women ages 50–74 with a mammogram in the past two years | 2022 | 79% | 77% | 86% | 14 of 51 |
| Percent of women ages 21–65 with a Pap smear in the past three years | 2022 | 85% | 82% | 89% | 7 of 51 |
| Percent of women ages 45–74 who received a sigmoidoscopy or a colonoscopy in the past 10 years or a fecal occult blood test in the past two years | 2022 | 61% | 60% | 69% | 34 of 51 |
| Percent of women with a recent live birth with a flu shot in the 12 months before delivery or during pregnancy | 2021 | 57% | 57% | 78% | 21 of 33 |
| Percent of women age 65 and older who ever received a pneumonia vaccine | 2022 | 70% | 73% | 80% | 43 of 51 |
| Percent of women ages 18–64 who have ever had an HIV or AIDS test | 2022 | 54% | 44% | 67% | 3 of 51 |
| Percent of women with a recent live birth who reported being asked about depression during a maternal postpartum checkup visit | 2021 | 84% | 89% | 96% | 29 of 33 |
| Percent of women with a recent live birth who had teeth cleaned during pregnancy by a dentist or dental hygienist | 2021 | 43% | 45% | 56% | 25 of 33 |

[a] Not all indicators available for all 50 states plus D.C. [b] U.S. rate is the 51-state median.