

# Tacoma, WA

Ranking Summary (of 306 Local Areas)	2016 Scorecard		2012 Scorecard (Rev.) <sup>a</sup>	
	Quintile	Rank	Quintile	Rank
<b>OVERALL</b>	<b>3</b>	<b>153</b>	<b>3</b>	<b>184</b>
Access & Affordability	2	122	4	199
Prevention & Treatment	4	223	5	262
Avoidable Hospital Use & Cost	2	78	2	86
Healthy Lives	3	170	3	149

Change in Performance <sup>c</sup>	2016 Scorecard	
	count	percent
Indicators with trends	33	100%
Area rate improved	15	45%
Area rate worsened	1	3%
Little or no change in area rate	17	52%

Percentage of Indicators This Area Ranked in the:	2016 Scorecard	
	count	percent
Total indicators	36	100%
Top 10th percentile	3	8%
Top quintile	8	22%
2nd quintile	10	28%
3rd quintile	7	19%
4th quintile	5	14%
Bottom quintile	6	17%
Bottom 10th percentile	3	8%

Demographic Characteristics <sup>b</sup>	Tacoma, WA	U.S. Average
<b>Total Population</b>	<b>807,790</b>	<b>313,922,028</b>
Median Household Income	\$62,732	\$58,489
Below 200% of Federal Poverty Level (FPL)	29%	34%
200% to 399% of FPL	32%	30%
400% of FPL or higher	39%	36%
% White Race, Non-Hispanic	69%	62%
% Black Race, Non-Hispanic	6%	12%
% Other Race, Non-Hispanic	15%	8%
% Hispanic Ethnicity	10%	17%

Estimated Impact of Local Improvement <sup>d</sup> If Tacoma, WA improved its performance to the level of the best-performing local area for this indicator, then:	
29,767	more adults (ages 19–64) would be covered by health insurance (public or private), and therefore would be more likely to receive health care when needed
101,943	more adults (age 18 and older) would have a usual source of care to help ensure that care is coordinated and accessible when needed
32,410	more adults would receive age and gender appropriate recommended preventive care, such as colon cancer screenings, mammograms, and pap smears
2,091	fewer Medicare beneficiaries would receive an unsafe medication
193	fewer premature deaths (before age 75) would occur from causes that are potentially treatable or preventable with timely and appropriate care
173	fewer hospital readmissions would occur among Medicare beneficiaries (age 65 and older)
1,884	fewer emergency department visits for nonemergent or primary-care-treatable conditions would occur among Medicare beneficiaries
40,413	fewer adults (ages 18–64) would have lost six or more teeth because of tooth decay, infection, or gum disease

## Notes

(a) Ranks from the 2012 edition of the Local Scorecard have been revised to match methodology and measure definitions used in the 2016 edition.

(b) The Local Scorecard reports at the level of the Hospital Referral Region (HRR), an area representing the local market for health care services, defined by health care utilization patterns. HRR boundaries may differ from other commonly used region definitions (e.g., county or Metropolitan Statistical Areas).

(c) Improved or worsened denotes a change of at least one-half (0.5) of a standard deviation (a statistical measure of variation) larger than the indicator's distribution among all Hospital Referral Regions (HRRs) over the two time points. Little or No change denotes no change in rate or a change of less than one-half of a standard deviation.

(d) The table shows the estimated impact if this HRR's performance improved to the rate of the best-performing HRR for eight Scorecard indicators. Benchmark HRRs, those with the best rate, have an estimated impact of zero (0).

[See the interactive 2016 Local Scorecard report and link to Scorecard methodology at www.CMWF.org.](#)

Table 1. Local Area Health System Performance Indicator Data by Dimension

Dimension and Indicator	Data Year	Tacoma, WA			Benchmarks			Data Year	Area Rate	U.S. Average	Substantial Change Over Time <sup>1</sup>	
		Performance Quintile	Rank (of 306 )	Area Rate	Washington Rate	U.S. Average	Best Area Rate					
ACCESS & AFFORDABILITY								2016 Scorecard			Baseline	
Adults ages 19–64 uninsured	2014	2	91	13%	13%	16%	4%	2012	19%	21%	Improved	
Children ages 0–18 uninsured	2014	1	22	4%	5%	6%	2%	2012	8%	7%	Improved	
Adults who went without care because of cost in past year	2013/14	3	170	16%	14%	15%	6%	2011/12	18%	15%	No Change	
At-risk adults without a routine doctor visit in past two years	2013/14	5	272	19%	18%	14%	6%	2011/12	22%	14%	Improved	
Adults without a dental visit in past year	2012 & 2014	2	85	14%	13%	15%	9%	--	--	--	--	
PREVENTION & TREATMENT								2016 Scorecard			Baseline	
Adults with a usual source of care	2013/14	5	259	73%	77%	79%	90%	2011/12	76%	79%	Worsened	
Adults with age and gender appropriate cancer screenings	2012 & 2014	2	99	69%	71%	70%	79%	--	--	--	--	
Adults with age appropriate vaccines	2013/14	1	36	42%	37%	35%	49%	2011/12	36%	35%	Improved	
Medicare beneficiaries who received at least one drug that should be avoided in the elderly	2013	4	199	18%	16%	17%	9%	2011	22%	20%	Improved	
Medicare beneficiaries with dementia, hip/pelvic fracture, or chronic renal failure who received a prescription drug that is contraindicated for that condition	2013	3	144	18%	15%	20%	9%	2011	19%	23%	No Change	
Risk-adjusted 30-day mortality among Medicare beneficiaries hospitalized for heart attack, heart failure, pneumonia, or stroke	07/2011 - 06/2014	5	266	13.5%	13.4%	12.8%	11.3%	07/2010 - 06/2013	14.4%	13.2%	Improved	
Hospital safety composite score <sup>2</sup>	07/2012 - 06/2014	5	292	1.0	0.9	0.8	0.6	07/2011 - 06/2013	1.2	0.9	Improved	
Hospitalized patients given information about what to do during their recovery at home	2014	1	24	89%	88%	86%	91%	2013	88%	86%	No Change	
Hospitalized patients who reported hospital staff always managed pain well, responded when needed help to get to bathroom or pressed call button, and explained medicines and side effects	2014	5	275	63%	66%	68%	76%	2013	61%	68%	Improved	
Home health patients who get better at walking or moving around	2014	4	220	59%	58%	63%	70%	2012	53%	59%	Improved	
Home health patients whose wounds improved or healed after an operation	2014	2	91	91%	88%	89%	96%	2012	92%	89%	No Change	
High-risk nursing home residents with pressure sores	1/2015 - 9/2015	1	56	5%	5%	6%	1%	1/2013 - 9/2013	6%	6%	Improved	
Long-stay nursing home residents with an antipsychotic medication	1/2015 - 9/2015	3	153	18%	17%	18%	7%	1/2013 - 9/2013	23%	21%	Improved	

Table 1. Local Area Health System Performance Indicator Data by Dimension (continued)

Dimension and Indicator	Data Year	Tacoma, WA			Benchmarks			Data Year	Area Rate	U.S. Average	Substantial Change Over Time <sup>1</sup>
		Performance Quintile	Rank (of 306 )	Area Rate	Washington Rate	U.S. Average	Best Area Rate				
2016 Scorecard								Baseline			
AVOIDABLE HOSPITAL USE & COST											
Hospital admissions among Medicare beneficiaries for ambulatory care–sensitive conditions, ages 65–74, per 1,000 beneficiaries	2014	2	72	21	17	27	9	2012	26	29	Improved
Hospital admissions among Medicare beneficiaries for ambulatory care–sensitive conditions, age 75 and older, per 1,000 beneficiaries	2014	2	74	55	46	66	33	2012	65	70	Improved
Medicare 30-day hospital readmissions, rate per 1,000 beneficiaries	2014	2	94	24	20	27	10	2012	31	34	Improved
Short-stay nursing home residents readmitted within 30 days of hospital discharge to nursing home	2012	2	77	18%	17%	20%	12%	2010	19%	22%	No Change
Long-stay nursing home residents hospitalized within a six-month period	2012	3	126	17%	13%	17%	5%	2010	18%	19%	No Change
Home health patients also enrolled in Medicare with a hospital admission	2014	1	9	14%	15%	16%	12%	2012	14%	17%	No Change
Potentially avoidable emergency department visits among Medicare beneficiaries, per 1,000 beneficiaries	2013	1	47	158	156	181	122	2011	147	185	No Change
Total reimbursements per enrollee (age 18-64) with employer-sponsored insurance	2014	4	240	\$4,994	\$4,357	\$4,569	\$2,720	2013	\$5,142	\$4,489	No Change
Total Medicare (Parts A & B) reimbursements per enrollee	2014	1	52	\$7,388	\$7,125	\$8,819	\$5,593	2012	\$7,480	\$8,854	No Change
2016 Scorecard								Baseline			
HEALTHY LIVES											
Mortality amenable to health care, deaths per 100,000 population	2012-13	2	104	75	62	84	47	2010-11	77	85	No Change
Breast cancer deaths per 100,000 female population	2012-13	3	123	22.0	20.8	22.8	12.3	2010-11	22.6	23.7	No Change
Colorectal cancer deaths per 100,000 population	2012-13	1	43	13.8	14.1	15.9	9.6	2010-11	16.7	16.7	Improved
Suicide deaths per 100,000 population	2012-13	4	231	16.9	14.7	12.9	6.0	2010-11	17.6	12.5	No Change
Infant mortality, deaths per 1,000 live births	2012-13	2	103	5.5	4.9	6.0	2.5	2010-11	4.9	6.1	No Change
Adults ages 18–64 who report fair/poor health or activity limitations because of physical, mental, or emotional problems	2013/14	5	270	33%	27%	25%	13%	2011/12	31%	25%	No Change
Adults who smoke	2013/14	4	188	21%	16%	19%	7%	2011/12	19%	19%	No Change
Adults ages 18–64 who are obese (BMI >= 30)	2013/14	3	179	33%	27%	28%	14%	2011/12	31%	28%	No Change
Adults ages 18–64 who have lost six or more teeth because of tooth decay, infection, or gum disease	2012 & 2014	3	156	12%	8%	10%	2%	--	--	--	--

Table 1 Notes:

\* Data for this area not available for this indicator.

-- Historical data not available or not comparable over time.

(1) Improved or worsened denotes a change of at least one-half (0.5) of a standard deviation (a statistical measure of variation) larger than the indicator's distribution among all Hospital Referral Regions (HRRs) over the two time points. No change denotes no change in rate or a change of less than one-half of a standard deviation.

(2) Values are the unweighted average of the region's hospitals' safety composite (PSI 90) score.